

目次

植物生理与育种

- 杜鹃花属植物种间杂交向性研究 庄平 (1281)
- 深挖垦复对锥栗重要农艺性状及土壤理化性质的影响 江锡兵, 杨龙, 吴聪连, 章平生, 龚榜初, 赖俊声 (1287)
- 模拟践踏对中位泥炭藓叶绿素荧光特性的影响 范贝贝, 卜兆君, 刘超, 胡雪凤, 陈永达 (1298)
- 两种砧木对樱桃番茄青枯病抗性及其根际微生物数量的影响 赵文宗, 郑旭阳, 张映卿, 钟川, 阳燕娟, 于文进 (1307)
- 蚕豆幼苗光合特性对土荆芥挥发性物质胁迫的响应 谷从璟, 周健, 马丹炜, 黄素, 孟巧巧 (1317)
- 大花蕙兰‘红酒’×莲瓣兰‘边草素花’杂交种组培快繁技术 刘洋, 王玉英, 李枝林, 李茹 (1327)
- 台湾早熟蜜桃优良芽变新种质‘五月红’的鉴定 潘介春, 龙嵩宇, 丁峰, 徐石兰, 黄幸, 黄思婕, 杨亚涵, 李峰, 张树伟 (1334)
- 基于生理生化指标选择红锥二代优良家系 谭长强, 申文辉, 刘秀, 曹艳云, 黄志玲, 郝海坤 (1342)
- 金铁锁转录因子 *ptMYC2* 的克隆和生物信息学分析 孟文俊, 张爱丽, 蒋乐晓, 朱文杰, 段丽, 钱子刚 (1350)

植物生态与地理

- 贵州特有植物红花瘤果茶种群空间格局及数量动态研究 郭米香, 杨乃坤, 刘海燕, 汤升虎, 范志伟, 邹天才 (1359)
- 北京市主城区城市森林景观格局特征分析 佟济宏, 王新杰, 汪锦, 傅锋 (1370)
- 贵州楼梯草属(荨麻科)植物种类组成及分布格局 崔兴勇, 安明志, 施金竹, 刘锋 (1379)
- 长柄双花木叶性状异速生长关系随发育阶段和海拔梯度的变化 耿梦娅, 陈芳清, 吕坤, 王玉兵, 官守鹏, 刘杨赞 (1387)
- 高盆樱桃与钟花樱桃的地理分布模拟及生态特征比较分析 朱淑霞, 朱弘, 程琳, 伊贤贵, 王贤荣 (1398)
- 苹果属, 东南亚大陆菊科一新归化属(英文) 沈佳豪, 牛恂豪, 张国进, 付志奎, INTA Angkhana, 高天刚 (1407)
- 广西禾本科植物二新记录属 陆昭岑, 邹春玉, 彭华 (1412)
- 中国苋科苋属新记录种——鲍氏苋和布氏苋 徐晗, 李振宇 (1416)
- 山东苔藓植物新记录 任昭杰, 田雅娴, 赵遵田 (1420)

责任编辑 何永艳 李莉 周翠鸣 蒋巧媛
 责任校对 周翠鸣 何永艳 周翠鸣 蒋巧媛
 英文编辑/校对 周翠鸣 李莉 何永艳 蒋巧媛
 封面/版式设计 李莉 何永艳 蒋巧媛 李莉

期刊基本参数: CN 45-1134/Q * 1981 * m * A4 * 144 * zh+en * P * ¥45.00 * 1200 * 18 * 2019-10

封面说明: 樱花烂漫几多时, 桃红柳绿两不知。阳春三月, 樱花盛开, 繁茂如云, 灿若朝霞。引来一大批游人驻足观赏, 甚是壮观。提起樱花, 人们脑海中大多浮现出花瓣如霏雪般, 婉转而下, 洒落满地的场景。樱花是许多品种的统称, 樱花品种的观赏特征各异, 使得园林景观更具观赏性。中国樱花品种的选育工作虽然起步较晚, 但中国是世界上拥有野生樱花最多的国家, 这一天然优势使得我国的樱花产业蒸蒸日上。
 照片示: 樱花属代表性植物。1. 市原虎之尾; 2. 海猫; 3. 菊枝垂樱; 4. 雨晴垂枝; 5. 尾叶樱桃; 6. 红笠; 7. 粉巨人; 8. 云南樱桃; 9. 八重红绯衣。(相关内容详见本期正文 1398~1406 页朱淑霞等的文章)

	1	2	3
	4		5 6
	7	8	9

CONTENTS

Plant Physiology and Breeding

- Direction of inter-specific cross of *Rhododendron* ZHUANG Ping (1281)
- Effects of deep digging and reclamation on important agronomic traits of *Castanea henryi* and soil physical and chemical properties JIANG Xibing, YANG Long, WU Conglian, ZHANG Pingsheng, GONG Bangchu, LAI Junsheng (1287)
- Effect of simulated trampling on the chlorophyll fluorescence characteristics of *Sphagnum magellanicum* FAN Beibei, BU Zhaojun, LIU Chao, HU Xuefeng, CHEN Yongda (1298)
- Effects of two rootstocks on bacterial wilt resistance and rhizospheric microorganisms of cherry tomato ZHAO Wenzong, ZHENG Xuyang, ZHANG Yingqing, ZHONG Chuan, YANG Yanjuan, YU Wenjin (1307)
- Response of photosynthetic characteristics of *Vicia faba* seedlings to volatile allelochemical stress from *Chenopodium ambrosioides* GU Congjing, ZHOU Jian, MA Danwei, HUANG Su, MENG Qiaoqiao (1317)
- Tissue culture and rapid propagation techniques in *Cymbidium hybridum* ‘hongjiu’ × *C. tortisepalum* ‘biancaosuhua’ LIU Yang, WANG Yuying, LI Zhilin, LI Ru (1327)
- Identification of a new bud sport germplasm ‘May Red’ from Taiwan precocious peach PAN Jiechun, LONG Qiangyu, DING Feng, XU Shilan, HUANG Xing, HUANG Sijie, YANG Yahan, LI Feng, ZHANG Shuwei (1334)
- Seedling selection of the second superior families of *Castanopsis hystrix* based on the analysis of physiological indices TAN Zhangqiang, SHEN Wenhui, LIU Xiu, CAO Yanyun, HUANG Zhiling, HAO Haikun (1342)
- Cloning and bioinformatics analysis of *Psammosilene tunicoides* transcription factor *ptMYC2* MENG Wenjun, ZHANG Aili, JIANG Lexiao, ZHU Wenjie, DUAN Li , QIAN Zigang (1350)

Plant Ecology and Geography

- Spatial distribution pattern and quantitative dynamics of the endemic plant *Camellia rubituberculata* in Guizhou Province GUO Mixiang, YANG Naikun, LIU Haiyan, TANG Shenghu, FAN Zhiwei, ZOU Tiancai (1359)
- Landscape pattern analysis of urban forest in central Beijing TONG Jihong, WANG Xinjie, WANG Jin, FU Feng (1370)
- Species composition and distribution pattern of *Elatostema* (Urticaceae) in Guizhou CUI Xingyong, AN Mingtai, SHI Jinzhu, LIU Feng (1379)
- Changes of allometric relationships among leaf traits of *Disanthus cercidifolius* var. *longipes* in different ontogenetic stages and altitude gradients GENG Mengya, CHEN Fangqing, LÜ Kun, WANG Yubing, GUAN Shoupeng, LIU Yangyun (1387)
- Modeling geographical distribution pattern and comparison of ecological characteristics between *Cerasus cerasoides* and *C. campanulata* ZHU Shuxia, ZHU Hong, Cheng Lin, YI Xiangui, WANG Xianrong (1398)
- Centratherum* Cass., a newly naturalized genus of family Asteraceae from mainland of Southeast Asia SHEN Jiahao, NIU Guohao, ZHANG Guojin, FU Zhixi, INTA Angkhana, GAO Tiangang (1407)
- Two newly recorded genera of Poaceae from Guangxi, China LU Zhaoen, ZOU Chunyu, PENG Hua (1412)
- Amaranthus powellii* S. Watson and *A. bouchonii* Thell., two newly naturalized species in China XU Han, LI Zhenyu (1416)
- New additions to the bryophyte flora of Shandong Province, China REN Zhaojie, TIAN Yaxian, ZHAO Zuntian (1420)

Cover images: Selective presentation of *Cerasus* plants. 1. *Cerasus serrulata* var. *lannesiana* ‘Lchihara’; 2. *Cerasus* ‘Umineko’; 3. *C. serrulata* var. *lannesiana* ‘Plena-pendula’; 4. *C. subhirtella* ‘Ujou-shidare’; 5. *C. dielsiana*; 6. *C. serrulata* var. *lannesiana* ‘Benigasa’; 7. *Cerasus* ‘Pink giant’; 8. *C. yunnanensis*; 9. *C. campanulata*. (For detail, please see the text by ZHU Shuxia et al. on page 1398–1406)

	1	2	3
			5 6
	4		9
	7	8	