

福建假卫矛属(卫矛科)—新种——德化假卫矛

黄志森¹, 林彦云²

(1. 戴云山国家级自然保护区, 福建 德化 362500; 2. 泉州市林业局, 福建 泉州 362000)

摘要: 对新种德化假卫矛进行描述, 该种仅见福建德化九仙山。

关键词: 假卫矛属; 德化假卫矛; 卫矛科; 新种; 中国

中图分类号: Q949.754.7 **文献标识码:** A **文章编号:** 1000-3142(2008)04-0458-02

Microtropis dehuaensis Z. S. Huang & Y. Y. Lin, a new species of Celastraceae from Fujian

HUANG Zhi-Sen¹, LIN Yan-Yun²

(1. Fujian Daiyunshan National Nature Reserve, Dehua 362500, China;

2. Forestry Bureau of Quanzhou City, Quanzhou 362000, China)

Abstract: A new species of the genus *Microtropis*, *M. dehuaensis* Z. S. Huang & Y. Y. Lin, is described from Jiuxianshan, Fujian.

Key words: *Microtropis*; *Microtropis dehuaensis* Z. S. Huang & Y. Y. Lin; Celastraceae; new species; China

德化假卫矛 新种 图版 I :A,B

***Microtropis dehuaensis* Z. S. Huang & Y. Y. Lin,
sp. nov. Plate I :A,B.**

Species proxima *M. fokienensis* Dunn, sed foliis oblongo-lanceolatis vel elliptico-lanceolatis, apice acuminatis vel abrupte caudatis, interdum caudato ipso saepe falcatis, inflorescentiis vulgo 3—4-dichotomis, pedunculis 1.8—2.8 cm longis, gracilibus, floribus 5-meris differt.

Frutex usque 3 m altus, omnis glaber, ramulis fusco-purpureis, hornotinis subgracilibus paulo angularibus annotinis vetustisque teretibus atropurpureis, lenticellis minutis pallidis, internodiis 1.5—2.5 cm longis. Folia opposita, chartacea oblongo-lanceolata vel elliptico-lanceolata, 5—9 cm longa, 1.5—2.8 cm lata, apice acuminata vel abrupte caudata interdum caudato ipso saepe falcata, basi late cuneata vel cuneata, margine integra paulo revoluta, supra saturate viridia, subtus griseo-viridia, cos-

ta utrinque conspicue elevata, nervis lateralibus utrinsecus 6—9, tenuibus indistinctis; petoli 5—8 mm longi. Inflorescentia axillaris thyrsoides vulgo 3—4-dichotoma 3—3.5 cm longa, 1.5—2.5 cm lata, pedunculis 1.8—2.8 cm longis, gracilibus, ramis primis c. 4 mm longis, secundis c. 3 mm longis, tertiiis necnon quartis sensim brevibus, pedicellis c. 2 mm longis, floribus albis 5-meris, sepalis semiorbiculatis c. 2 mm latis, petalis oblongo-ellipticis c. 2.5 mm longis, staminibus pistillo longioribus, antheris oblongo-cordatis, ovario anguste conoideo. Capsulae ellipsoideae atro-nigrescentes c. 1.2 cm longae, prope media crassior 2—3 mm diam. a medio sursum ad apicem sensim tenues, apice rostriformes, semine elliptico c. 7.5 mm longo fulvo.

China. Fujian(福建); Dehua(德化), Jiuxianshan(九仙山), alt. 1 000—1 200 m, in mixed forest, 2006-11-22, Z. S. Huang & Y. Y. Lin(黄志森、林彦云)0624(IBK); the same locality, 2007-08-14, Z. S.


收稿日期: 2007-10-09 修回日期: 2007-12-29

作者简介: 黄志森(1966-), 男, 福建德化县人, 高级工程师, 从事福建戴云山国家级自然保护区保护管理工作, E-mail: dysbhq@dhca.net.

Huang & Y. Y. Lin 0778(fr.) (holotype, IBK; isotype, Fujian Daiyunshan National Nature Reserve;

isotype, Forestry Bureau of Quanzhou City).

灌木,高可达3 m,全部无毛;小枝暗紫色,当年



图版 I 德化假卫矛果期生境照片 A. 植物外形; B. 果枝。
Plate I *Microtropis dehuaensis* Z. S. Huang & Y. Y. Lin. A. habit; B. fruiting branch.

生枝纤细,略有棱,一年生枝、老枝圆柱状,暗紫色,散生细小灰白色皮孔,节间长1.5~2.5 cm。叶对生,纸质,长圆状披针形或椭圆状披针形,长5~9 cm,宽1.5~2.8 cm,先端渐尖或尾尖,有时镰状弯向一侧,基部阔楔形或楔形,边缘全缘稍反卷,叶面深绿色,背面灰绿色,中脉在两面明显凸起,侧脉每边6~9条,纤细,不明显;叶柄长5~8 mm。花序腋生,聚伞花序

通常3~4次二岐分枝,长3~3.5 cm,宽1.5~2.5 cm,花序梗长1.8~2.8 cm,纤细,第一次分枝长约4 mm,第二次分枝长约3 mm,第三、第四次分枝逐渐渐短,小花梗长约2 mm,花白色,5数,萼片半圆形,宽约2 mm,花瓣长圆状椭圆形,长约2.5 mm,雄蕊比雌蕊长,花药长圆状心形,子房狭圆锥状。蒴果椭圆状,(下转第538页 Continue on page 538)

物,避开银胶菊的化感作用武器。如果茶条木不受银胶菊的化感作用影响,又作为重要的能源植物,在银胶菊入侵严重,适宜茶条木生长的地区,如广西百色地区大量种植茶条木,影响银胶菊生长的物理环境如光照、生长空间等,控制其生长和繁殖,既有效防治银胶菊,也为能源植物茶条木的推广种植奠定基础。此外,探索银胶菊的化感作用,用其化感成分开发除草剂,充分利用的同时达到防治的效果。

参考文献:

- Adkins SW, Sowerby MS. 1996. Allelopathic potential of the weed, *Parthenium hysterophorus* in Australia[J]. *Plant Protection Quarterly*, 11: 20—23
- Batish DR, Singh HP, Kohli RK, et al. 2002. Allelopathic effects of parthenin against two weedy species, *Avena fatua* and *Bidens pilosa*[J]. *Environ Experimental Bot*, 47: 149—155
- Chippendale JF, Panetta FD. 1994. The cost of *Parthenium hysterophorus* in the Queensland cattle industry[J]. *Plant Protection Quarterly*, 9: 73—76
- Hsu LM, Chiang MY. 2004. Seed germination and chemical control of Parthenium weed(*Parthenium hysterophorus*)[J]. *Weed Sci Bull*, 25(1): 11—21
- Kanchan SD, Jayachandra. 1979. Allelopathic effects of *Parthenium hysterophorus* I. Exudation of inhibitors through roots[J]. *Plant Soil*, 53: 27—35
- Kanchan SD, Jayachandra. 1980. Allelopathic effects of *Parthenium hysterophorus* I. Identification of inhibitors[J]. *Plant Soil*, 55: 67—75
- Kohli RK, Batish DR. 1994. Exhibition of allelopathy by *Parthenium hysterophorus* in agroecosystems[J]. *Trop Ecol*, 35: 295—307
- More PR, Vadlamudi VP, Qureshi MI. 1982. Note on the toxicity of *Parthenium hysterophorus* in livestock[J]. *Indian J Animal Sci*, 52: 456—457
- Pandey DK. 1994. Inhibition of *Salvinia* (*Salvinia molesta* Mitch-
- ell) by *Parthenium* (*Parthenium hysterophorus*). II. Relative effect of flower, leaf, stem, and root residue on *salvinia* and *padhy*[J]. *J Chem Ecol*, 20(12): 3 123—3 131
- Pandey DK. 1996. Phytotoxicity of sesquiterpene lactone parthenin on aquatic weeds[J]. *J Chem Ecol*, 22: 151—160
- Ramaswami PP. 1997. Proc. First International Conference on Parthenium Management, 1: 77—80
- Singh HP, Batish DR, Kohli RK. 2002. Effect of Parthenin-A sesquiterpene lactone from *Parthenium hysterophorus*—on early growth and physiology of *Ageratum conyzoides* [J]. *J Chem Ecol*, 28(11): 2 169—2 179
- Singh HP, Batish DR, Pander JK, et al. 2005. Phytotoxic effects of *Parthenium hysterophorus* residues on three *Brassica* species [J]. *Weed Biology and Management*, 5(3): 105—109
- Tamado T, Milberg P. 2000. Weed flora in arable fields of eastern Ethiopia with emphasis on the occurrence of *Parthenium hysterophorus*[J]. *Weed Res*, 40: 507—521
- Tang SC(唐赛春), Lü SH(吕仕洪), He CX(何成新), et al. 2008. Distribution and harmful effects of alien invasive plant *Parthenium hysterophorus* in Guangxi(外来入侵植物银胶菊在广西的分布与危害)[J]. *Guizhou Botanical Research*(广西植物), 28(2): 197—200
- Tudor GD, Ford AL, Armstrong TR, et al. 1982. Taints in meat from sheep grazing *Parthenium hysterophorus*[J]. *Australian J Experimental Agric Animal Husbandry*, 22: 43—46
- Wang KM(王康满), Hou YT(侯元同). 2004. *Parthenium*(Asteraceae) a newly naturalized record genus in Shandong Province(山东归化植物—新记录属—银胶菊属)[J]. *J Qufu Norm Univ*(曲阜师范大学学报), 30(1): 83—84
- Williamson GB, Richardson D. 1988. Bioassays for allelopathy: Mersuring treatment responses with independent controls[J]. *J Chem Ecol*, 14(1): 181—187
- Zhu SX(朱世新), Qin HN(覃海宁), Chen YL(陈艺林). 2005. Alien species of Compositae in China(中国菊科植物外来种概述)[J]. *Guizhou Botanical Research*(广西植物), 25(1): 69—76

(上接第 459 页 Continue from page 459)

黑紫色,长约 1.2 cm,近中部最粗,径 2~3 mm,从中部向顶端渐细,顶端喙状,种子椭圆状,长约 7.5 mm,黄褐色。

本种与福建假卫矛相近,但叶长圆状披针形或椭圆状披针形,先端渐尖或尾状,有时镰状弯向一侧,花序 3~4 次二歧分枝,花序梗长 1.8~2.8 cm,纤细,花 5 数,易于区别。

致谢 在本种鉴定过程中,得到福建省亚热带植物研究所张永田研究员的悉心指导,谨致谢忱。

参考文献:

- 诚静容,高作经. 1999. 卫矛科. 中国植物志[M]. 北京:科学出版社,45(3): 150—174
- Kao TC(高作经), Cheng CY(诚静容). 1988. New Taxa of the Chinese Celastraceae (1)(中国卫矛科新分类群(1))[J]. *Acta Phytotax Sin*(植物分类学报), 26(4): 310—314
- Shui YM(税玉民), Chen WH(陈文红). 2002. A new species of the genus *Microtropis*(Celastraceae) from SE Yunnan, China(滇东南假卫矛属—新种)[J]. *Acta Bot Yunnan*(云南植物研究), 24(6): 707—708