

瓣片顶端 2 微裂或 2 浅裂。雌蕊群由 3 心皮组成。

2 种, 隶属于 2 系。分布于四川和新疆。

系 1. 征镒翠雀花系

Ser. **Flavidilimba** W. T. Wang, ser. nov. Type:

Delphinium zhenyianum Y. Luo & Q. E. Yang.

Flores parvi, in racemum compositum dispositi. Sepala 5–7 mm longa, prope apicem sub angulo 45° incurvata. Carpella glabra.

花小, 组成复总状花序。萼片长 5~7 mm, 近顶端成直角向内弯曲。心皮无毛。

1 种, 产于四川西部。

160. 征镒翠雀花

Delphinium zhengyianum Y. Luo & Q. E.

Yang in herb., sp. nov. Type 四川 (Sichuan): 道孚县, 各卡乡, 各卡村 (Daofu County, Geka Town, Geka Village), alt. 3 300 m, 沟边, 花白色, 上萼片深紫色 (by stream, fls. white, upper sepal deep purple), 2001-08-02, 罗艳 (Y. Luo) 351 (holotype, PE); 雅江县, 剪子弯兵站 (Yajiang County, Jianziuan military depot), alt. 3 200 m, 路边草丛中, 萼片白色, 上萼片深紫色 (among grasses by road, sepals white, upper sepal deep purple), 2001-08-29, 罗艳 357 (paratype, PE)。

Herbs perennial, ca. 1 m tall. Stems below very sparsely spreading-pubescent, above subglabrous, simple, several-leaved. Basal leaves not seen. Cauline leaves long or shortly petiolate; blades papery, pentagonal, 6–7 × 8–11 cm, at base cordate, 3-parted, central primary lobe rhombic, near the middle 3-lobed, secondary lobes with a few lobules and teeth, lateral primary lobes obliquely flabellate, unequally 2-parted; surfaces both densely puberulous; petioles 4.6–9 cm long. Compound racemes terminal, ca. 32 cm long;

bracts foliaceous; pedicels 5–13 mm long, spreading-puberulous, below to above 2-bracteolate; bracteoles filiform, 3–5 mm long, glabrous. Sepals outside puberulous, inside subglabrous, near apex by right angle incurved, upper sepal deep purple, navicular-ovate, 6 × 2.2 mm, spur subulate, 6–7 mm long, at base 1.5–2 mm across, lateral sepals white, oblong, 6 × 2 mm, lower sepals also white, navicular-oblong, 6–7 × 2 mm. Petals yellowish, glabrous, undivided. Staminodes concolour with petals; limbs ovate, 3 × 2 mm, 2-lobed, above ciliate, adaxially on the middle white-barbate; claws linear, 2.8 mm long, glabrous. Stamens glabrous. Carpels 3, glabrous.

特产于四川西部。生于山地沟边或路边草丛中, 海拔 3 200~3 300 m。

系 2. 文采新翠雀花系

Ser. **Flavidilimboida** W. T. Wang, ser.

nov. Type: *Delphinium neowentsaii* C. Y. Yang.

Flores majores, in racemum simplicem dispositi. Sepala ca. 12 mm long, non incurvata. Carpellorum ovaria dense puberula.

花较大, 组成简单总状花序。萼片长约 12 mm, 不向内弯曲。心皮子房密被短柔毛。

1 种, 产于新疆乌鲁木齐县。

161. 文采新翠雀花

Delphinium neowentsaii C. Y. Yang in Acta Phytotax. Sin. 43(1): 68, fig. 1. 2005. Type 新疆: 乌鲁木齐县, 天山, 水西沟, alt. 2 750 m, 林缘, 2001-08-18, 杨宗宗 0431 (holotype, XJA; isotype, PE)。

草本高约 30 cm。茎下部密被反曲糙伏毛, 中部以上被短柔毛或无毛, 等距生叶。基生叶具长柄; 叶片五角形, 约 4 × 7 cm, 基部深心形, 3 深裂, 中央深裂片宽菱形, 在中部 3 裂, 2 回裂片